

Student Name.....Student Number.....Section.....

You may use the following information wherever necessary:

a)  $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1} = 0.0821 \text{ atm dm}^3 \text{ mol}^{-1} \text{ K}^{-1} = 0.0821 \text{ atm L mol}^{-1} \text{ K}^{-1}$

b)  $k = Ae^{-E_a/RT}$

g)  $K_w = 1.0 \times 10^{-14}$  at 298 K

c)  $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \times \frac{(T_2 - T_1)}{T_1 T_2}$

h)  $K_a(\text{CH}_3\text{COOH}) = 1.8 \times 10^{-5}$

d)  $\ln \frac{[A]_t}{[A]_0} = -kt$

i)  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

e)  $t_{1/2} = \frac{\ln 2}{k} = \frac{0.693}{k}$

j)  $\text{pH} = \text{pK}_a + \log \frac{[\text{base}]}{[\text{acid}]}$

f)  $K_p = K_c (0.0821T)^{\Delta n(\text{gas})}$

k)  $E = E^\circ - \frac{0.059}{n} \log Q$

**Section A: Multiple Choice**

Select the best answer for each question and shade the letter corresponding to the answer on the answer sheet provided. [35 marks]

**Questions 1-3**The reaction  $2\text{NO}_2^-(\text{aq}) + 4\text{H}^+ + 2\text{I}^-(\text{aq}) \rightarrow \text{I}_2 + 2\text{NO}(\text{g}) + 2\text{H}_2\text{O}(\text{l})$  is first order in nitrite ion and iodide ion and second order in hydrogen ion.

1. The rate law for the reaction is

A  $\text{Rate} = k [\text{NO}_2^-]^2 [\text{H}^+] [\text{I}]^2$

B  $\text{Rate} = k [\text{NO}_2^-]^2 [\text{H}^+] [\text{I}]$

C  $\text{Rate} = k [\text{NO}_2^-] [\text{H}^+]^2 [\text{I}]$

D  $\text{Rate} = k [\text{NO}_2^-] [\text{H}^+] [\text{I}]^2$

E  $\text{Rate} = k [\text{NO}_2^-]^2 [\text{H}^+]^4 [\text{I}]^2$

2. If the rate of the reaction is expressed in  $\text{M s}^{-1}$ , the correct unit for the rate constant,  $k$ , is

A  $\text{M}^{-2} \text{s}^{-1}$

B  $\text{M}^2 \text{s}^{-1}$

C  $\text{M s}^{-1}$

D  $\text{M}^{-2} \text{s}^{-2}$

E  $\text{M}^{-3} \text{s}^{-1}$

3. By what factor would the rate of the reaction change if the concentrations of all the reactants are doubled?

A 1/2

B 2


C 4

D 8

E 16

Student Name.....Student Number.....Section.....

4. Ammonia can be oxidized according to the equation:




If in a particular reaction the  $\Delta[\text{NO}]$  is  $0.006 \text{ mol dm}^{-3}$ , then  $\Delta[\text{O}_2]$ , in  $\text{mol dm}^{-3}$ , is

- A  $-5/4 \times 0.006$
- B  $5/4 \times 0.006$
- C  $-4/5 \times 0.006$
- D  $4/5 \times 0.006$
- E  $4 \times 5 \times 0.006$
5. Which statement best explains the observation that reaction rates increase when temperature is increased?
- A At a higher temperature the energy of activation is reduced.
- B At a higher temperature the energy of activation is increased.
- C At a higher temperature the concentration of the reactants is higher.
- D At a higher temperature a larger fraction of reactant molecules have sufficient energy to form the transition state.
- E At a higher temperature there is no need to form the transition state.
6. Which statement about catalysts is **NOT** true?
- A A catalyst has no effect on the enthalpy change for the reaction which it catalyses.
- B A catalyst does not participate in the reaction which it catalyses.
- C Catalysts are specific in their action.
- D A catalyst changes the rate of the forward and reverse reactions for a reversible reaction by the same factor.
- E A catalyst does not affect equilibrium position for a reversible reaction.
7. The following mechanism has been proposed for a reaction:
- Step 1:  $\text{H}_2\text{O}_2(\text{aq}) + \text{I}^-(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l}) + \text{IO}^-(\text{aq})$  slow
- Step 2:  $\text{IO}^-(\text{aq}) + \text{H}_2\text{O}_2(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g}) + \text{I}^-(\text{aq})$  fast
- Which statement is **NOT** consistent with this proposed mechanism?
- A The overall reaction is:  $2 \text{H}_2\text{O}_2(\text{aq}) \rightarrow 2 \text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$
- B  $\text{IO}^-$  is a reactive intermediate.
- C  $\text{I}^-$  is a catalyst.
- D The reaction is first order with respect to the catalyst.
- E The reaction is second order with respect to  $\text{H}_2\text{O}_2$ .

Student Name.....Student Number.....Section.....

8. 0.24 mol of  $\text{NO}_2$  and 0.20 mol of  $\text{Cl}_2$  were introduced into a  $1 \text{ dm}^3$  vessel at constant temperature. When the system reached equilibrium, 0.16 mol of  $\text{NOCl}$  was present.



Which set of values shows the concentration of each gas at equilibrium?

	$[\text{NO}_2]/\text{mol dm}^{-3}$	$[\text{Cl}_2]/\text{mol dm}^{-3}$	$[\text{NOCl}]/\text{mol dm}^{-3}$
A	0.08	0.12	0.16
B	0.08	0.04	0.16
C	0.08	0.08	0.16
D	0.16	0.08	0.16
E	0.12	0.12	0.16

9. The equilibrium constant for the reaction  $\text{P}(\text{aq}) \rightleftharpoons \text{Q}(\text{aq})$  is  $3.2 \times 10^{-5}$ .

Which of the following statements is **TRUE**?

- A The equilibrium concentration of P is less than that of Q.  
 B The equilibrium concentration of P is greater than that of Q.  
 C Adding a suitable catalyst will increase the equilibrium concentration of Q.  
 D Adding a catalyst will increase the value of the equilibrium constant.  
 E Adding more P to an equilibrium mixture of P and Q will increase the value of the equilibrium constant.
10. For which equilibrium system, at constant temperature, will decreasing the volume **not** cause the equilibrium position to shift?
- A  $2 \text{CO}(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2 \text{CO}_2(\text{g})$   
 B  $\text{C}(\text{s}) + \text{O}_2(\text{g}) \rightleftharpoons \text{CO}_2(\text{g})$   
 C  $\text{COCl}_2(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + \text{Cl}_2(\text{g})$   
 D  $2 \text{NH}_3(\text{g}) \rightleftharpoons \text{N}_2(\text{g}) + 3 \text{H}_2(\text{g})$   
 E  $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$

11. Consider the process:  $\text{Fe}_2\text{O}_3(\text{s}) + 3 \text{H}_2(\text{g}) \rightleftharpoons 3\text{H}_2\text{O}(\text{g}) + 2 \text{Fe}(\text{s}) \quad \Delta H = + 98.7 \text{ kJ}$

Which statement is **NOT** true for this system?

- A  $K_p = K_c$  at a stated temperature.  
 B Addition of some  $\text{H}_2$  to an equilibrium mixture will cause equilibrium to shift to the right.  
 C Increasing the mass of  $\text{Fe}_2\text{O}_3$  will cause equilibrium to shift to the right.  
 D The value of  $K_p$  can be increased by increasing the temperature.  
 E Decreasing the volume of the container does not upset equilibrium.

Student Name.....Student Number.....Section.....

12. For the reaction  $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$ ,  $K_p = 1.7$  at 298K. Five systems were set up with the initial partial pressure of each gas as shown in the table. In which system would the **forward** reaction occur to establish equilibrium?

	$p_i\text{PCl}_5/\text{atm}$	$p_i\text{PCl}_3/\text{atm}$	$p_i\text{Cl}_2/\text{atm}$
A	1	2	1
B	2	2	2
C	1	1	2
D	2	2	3
E	3	2	2

13. According to the Bronsted-Lowry definition, a **base** is a species which

- A donates a hydrogen atom.
- B donates a hydrogen ion.
- C accepts a hydrogen atom.
- D accepts a hydrogen ion.
- E accepts a hydroxide ion.

14. Which does **NOT** constitute an acid/base conjugate pair?

- A  $\text{H}_2\text{CO}_3/\text{HCO}_3^-$
- B  $\text{NH}_3/\text{NH}_2^-$
- C  $\text{NH}_4^+/\text{NH}_3$
- D  $\text{H}_3\text{O}^+/\text{OH}^-$
- E  $\text{HNO}_2/\text{NO}_2^-$

15. Which is a weak acid?

- A HI
- B  $\text{HClO}_4$
- C HBr
- D HF
- E HCl

16. Which set shows the substances in order of **increasing** acid strength?

- A  $\text{HClO}$ ,  $\text{HClO}_2$ ,  $\text{HClO}_3$ ,
- B  $\text{H}_2\text{SO}_4$ ,  $\text{H}_2\text{SO}_3$ ,  $\text{HSO}_4^-$
- C HCl, HBr, HF
- D HF,  $\text{H}_2\text{O}$ ,  $\text{NH}_3$
- E  $\text{HPO}_4^{2-}$ ,  $\text{H}_3\text{PO}_4$ ,  $\text{H}_2\text{PO}_4^-$ ,

Student Name.....Student Number.....Section.....

17. The acidity constant for an acid, HA, is  $2.5 \times 10^{-5}$ . The  $pK_b$  of its conjugate base is closest to
- A 4.6
  - B 9.4
  - C  $4.0 \times 10^{-10}$
  - D  $1.0 \times 10^{-14}$
  - E 14
18. Assuming all of the following solutions have the same molar concentration, which one would be expected to have the **lowest** pH?
- A  $FeCl_3$
  - B  $FeCl_2$
  - C  $CaCl_2$
  - D KCl
  - E  $BaCl_2$

---

**Questions 19-23** refer to the following titrations:

- A The titration of  $20.0 \text{ cm}^3$  of 0.1M HCl with 0.1 M NaOH
- B The titration of  $20.0 \text{ cm}^3$  of 0.1M HCl with 0.1 M  $NH_3$
- C The titration of  $20.0 \text{ cm}^3$  of 0.1M  $CH_3COOH$  with 0.1 M NaOH
- D The titration of  $20.0 \text{ cm}^3$  of 0.1M KOH with 0.1 M HCl
- E The titration of  $20.0 \text{ cm}^3$  of 0.1M  $HNO_3$  with 0.1 M KOH

For which titration

- 19. would there be a decreases in pH as the titrant is added?
- 20. would the pH be greater than 7 at the equivalence point?
- 21. would the pH be lower than 7 at the equivalence point?
- 22. would phenolphthalein (pH range 8.3 – 10.0) be unsuitable as an indicator?
- 23. would bromocresol green (pH range 3.8 – 5.4) be unsuitable as an indicator?

---

24. In which compound does hydrogen carry an oxidation number of  $-1$ ?

- A  $NH_4NO_3$
- B LiH
- C  $H_2O_2$
- D  $NaHSO_4$
- E HF

Student Name.....Student Number.....Section.....

25. In which compound does oxygen carry an oxidation number of -1?

- A  $\text{NaHSO}_4$
- B  $\text{NH}_4\text{NO}_3$
- C  $\text{H}_2\text{O}_2$
- D  $\text{Fe}_2\text{O}_3$
- E  $\text{FeO}$

26. In which compound does oxygen carry an oxidation number of +2?

- A  $\text{F}_2\text{O}$
- B  $\text{NH}_4\text{NO}_3$
- C  $\text{KHSO}_4$
- D  $\text{CuO}$
- E  $\text{Cu}_2\text{O}$

27. Which is **NOT** a redox reaction?

- A  $\text{Zn} + \text{H}_2\text{SO}_4 \rightarrow \text{ZnSO}_4 + \text{H}_2$
- B  $\text{Cu} + 2\text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 + \text{SO}_2 + 2\text{H}_2\text{O}$
- C  $2\text{NBr}_3 + \text{H}_2\text{O} \rightarrow \text{N}_2 + 4\text{Br}^- + 2\text{HOBr}$
- D  $\text{ZnCO}_3 \rightarrow \text{ZnO} + \text{CO}_2$
- E  $\text{XeF}_2 + 2\text{Cl}^- \rightarrow \text{Xe} + 2\text{F}^- + \text{Cl}_2$

28. Which is a disproportionation reaction?

- A  $\text{SO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_3$
- B  $\text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$
- C  $3\text{NO}_2 + \text{H}_2\text{O} \rightarrow 2\text{HNO}_3 + \text{NO}$
- D  $2\text{KMnO}_4 + 5\text{SO}_2 + 2\text{H}_2\text{O} \rightarrow 2\text{MnSO}_4 + \text{K}_2\text{SO}_4 + 2\text{H}_2\text{SO}_4$
- E  $\text{S}_2\text{O}_8^{2-} + 2\text{I}^- \rightarrow 2\text{SO}_4^{2-} + \text{I}_2$


29. Which quantities are conserved in a redox reaction?

- A Mass only.
- B Charge only.
- C Oxidation number.
- D Neither mass nor charge.
- E Both mass and charge.

Student Name.....Student Number.....Section.....

30. The e.m.f. of the cell: Pt (s)| H<sub>2</sub> (g)|HCl (aq)|| CuSO<sub>4</sub> (aq)| Cu (s) does **NOT** depend on
- A temperature.
  - B the size of the copper electrode.
  - C the concentration of HCl.
  - D the concentration of CuSO<sub>4</sub>.
  - E the pressure of H<sub>2</sub>.
31. When the contents of an electrochemical cell are at equilibrium, the e.m.f. of the cell
- A is zero.
  - B is at a maximum.
  - C is negative.
  - D is positive.
  - E cannot be measured.
- 

**Questions 32 - 35** concern the following graphs:




Select, from A to E, the graph which best represents:

32. Rate of reaction versus concentration of X for a reaction which is zero order in X.
33. Rate of reaction versus concentration of X for a reaction which is first order in X.
34. Rate of reaction versus time for a reversible process which attains equilibrium after some time.
35. The titration curve for the titration of a base with an acid.

Student Name.....Student Number.....Section.....

**SECTION B:** Answer **ALL** questions **in the spaces provided on the question paper.****Remember to include units in your answers wherever appropriate.**

1. The following data were obtained for the reaction:



Experiment	Initial [I <sup>-</sup> ]/M	Initial [BrO <sub>3</sub> <sup>-</sup> ]/M	Initial [H <sup>+</sup> ]/M	Initial Rate of I <sub>2</sub> formation/Ms <sup>-1</sup>
1	0.0020	0.0080	0.020	$8.89 \times 10^{-5}$
2	0.0040	0.0080	0.020	$1.78 \times 10^{-4}$
3	0.0020	0.0160	0.020	$1.78 \times 10^{-4}$
4	0.0020	0.0080	0.040	$3.56 \times 10^{-4}$

- a) Derive the rate law for the reaction. [3 ]

- b) i) Use the data from experiment 1 to find the value of the rate constant,
- $k$
- ,
- stating its correct units.
- [2]

- ii) What would be the value of the
- rate constant**
- if the concentration of all reactants were doubled? [1]

- c) What effect, if any, would doubling the concentration of the reactants have on the energy of activation for the process? [1]



Student Name.....Student Number.....Section.....




- d) What effect, if any, would increasing the temperature of the reaction mixture have on the energy of activation for the process? [1]
- e) What effect, if any, would increasing the temperature of the reaction mixture have on the value of the rate constant for the process? [1]
- f) What effect, if any, would using a catalyst have on the energy of activation for the process? [1]
2. The activation energy for the reaction:  $2 \text{N}_2\text{O} (\text{g}) \rightarrow 2 \text{N}_2 (\text{g}) + \text{O}_2 (\text{g})$  is  $200 \text{ kJ mol}^{-1}$ .  
How many times faster would this reaction proceed at  $230^\circ\text{C}$  than at  $200^\circ\text{C}$ ? [4]
3. The first order rate constant for the decomposition of a certain hormone in water at  $25^\circ\text{C}$  is  $0.0342 \text{ day}^{-1}$ .
- a) If a  $0.0200 \text{ M}$  solution of the hormone is stored for 40 days, what will be its concentration at the end of that period? [3]

Student Name.....Student Number.....Section.....

b) What is the half life of the hormone? [2]

c) How many days will it take for a sample of the hormone to be 65% decomposed? [2]

4. Use the given  $K_p$  values for the processes X and Y to find  $K_p$  for the process Z. [2]



Student Name.....Student Number.....Section.....

5. The equilibrium constant,  $K_p$ , for the dissociation of dinitrogen tetroxide to nitrogen dioxide is 11 at 398K. The reaction is:  $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ .

a) Find the equilibrium partial pressure of each gas when  $\text{N}_2\text{O}_4$ , at an initial pressure of 1.20 atm, dissociates at 398 K. [6]

b) Find the total pressure of the system at equilibrium. [1 ]

c) Find the percent dissociation of dinitrogen tetroxide. [1]

6. Find the pH of

a) 0.020 M NaOH [1]

Student Name.....Student Number.....Section.....

b) 0.020 M  $\text{CH}_3\text{COOH}$  [5]


c) a mixture of 20.0  $\text{cm}^3$  of 0.020 M  $\text{CH}_3\text{COOH}$  + 20.0  $\text{cm}^3$  of 0.020 M  $\text{NaOH}$  [7]

Student Name.....Student Number.....Section.....

d) a mixture of 30.0 cm<sup>3</sup> of 0.020 M CH<sub>3</sub>COOH + 20.0 cm<sup>3</sup> of 0.020 M NaOH [5]

e) a mixture of 20.0 cm<sup>3</sup> of 0.020 M CH<sub>3</sub>COOH + 30.0 cm<sup>3</sup> of 0.020 M NaOH [4]

7. Derive a balanced **ionic** equation for the reaction by writing half equations and then combining them.




Student Name.....Student Number.....Section.....

8. Use the following table of standard redox potentials wherever necessary.

	$E^{\circ}/V$
$MnO_4^- (aq) + 8 H^+ (aq) + 5 e^- \rightarrow Mn^{2+} (aq) + 4 H_2O (l)$	+1.51
$Cl_2 (g) + 2e^- \rightarrow 2Cl^- (aq)$	+1.36
$Ag^+ (aq) + e^- \rightarrow Ag (s)$	+0.80
$Fe^{3+} (aq) + e^- \rightarrow Fe^{2+} (aq)$	+0.77
$NiO_2 (s) + 2 H_2O (l) + 2e^- \rightarrow Ni(OH)_2 (s) + 2 OH^- (aq)$	+0.49
$Cu^{2+} (aq) + 2e^- \rightarrow Cu (s)$	+0.34
$2H^+ (aq) + 2e^- \rightarrow H_2 (g)$	0.00
$Ni^{2+} (aq) + 2e^- \rightarrow Ni (s)$	-0.25
$Fe^{3+} (aq) + 3e^- \rightarrow Fe (s)$	-0.036
$Cd^{2+} (aq) + 2e^- \rightarrow Cd (s)$	-0.40
$Fe^{2+} (aq) + 2e^- \rightarrow Fe (s)$	-0.44
$Cd(OH)_2(s) + 2e^- \rightarrow Cd (s) + 2 OH^- (aq)$	-0.81
$Mg^{2+} (aq) + 2e^- \rightarrow Mg (s)$	-2.38

- a) Rechargeable nickel-cadmium cells are used in calculators and other battery powered devices. The cell reaction is:



What is the cell potential of a standard nickel-cadmium cell? [1]

- b) The cell notation represents a **standard** galvanic cell:



- i) Write a balanced **ionic** equation for the cell reaction. [1]

- ii) Draw a **fully labeled** diagram of the galvanic cell. Show the direction of flow of electrons, the polarity of the electrodes and the concentration of all solutions. [5]

Student Name.....Student Number.....Section.....

- c) i) Find the standard cell potential for a cell in which the reaction:  
$$\text{Ag}^+(\text{aq}) + \text{Fe}^{2+}(\text{aq}) \rightleftharpoons \text{Ag}(\text{s}) + \text{Fe}^{3+}(\text{aq})$$
 takes place. [1]
- ii) Find  $K_c$  for the process:  $\text{Ag}^+(\text{aq}) + \text{Fe}^{2+}(\text{aq}) \rightleftharpoons \text{Ag}(\text{s}) + \text{Fe}^{3+}(\text{aq})$  [4]
- d) Find the e.m.f of the cell:  $\text{Cu}(\text{s}) | \text{Cu}^{2+}(0.001 \text{ M}) || \text{Cu}^{2+}(0.250 \text{ M}) | \text{Cu}(\text{s})$  [4]
- e) Explain why hydrochloric acid cannot be used to provide an acid medium with potassium manganate (VII) as an oxidizing agent. [2]

**END OF EXAMINATION**